



The Mechanisms by Which the Ketone Body D-β-Hydroxybutyrate May Improve the Multiple Cellular Pathologies of Parkinson's Disease

Nicholas G. Norwitz^{1*}, Michele T. Hu² and Kieran Clarke¹

¹ Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, United Kingdom, ² Nuffield Department of Clinical Neurosciences, Oxford University Hospitals NHS Foundation Trust, Oxford, United Kingdom

OPEN ACCESS

Edited by:

Massimo Lucarini,
Council for Agricultural and
Economics Research, Italy

Reviewed by:

Juan Segura-Aguilar,
Universidad de Chile, Chile

Bingxu Huang,
Jilin University, China
Marcelo M. S. Lima,

Federal University of Paraná, Brazil

*Correspondence:

Nicholas G. Norwitz
nicholas.norwitz@dpag.ox.ac.uk

Specialty section:

This article was submitted to
Nutrition and Food Science
Technology,
a section of the journal
Frontiers in Nutrition

Received: 23 January 2019

Accepted: 23 April 2019

Published: 14 May 2019

Citation:

Norwitz NG, Hu MT and Clarke K
(2019) The Mechanisms by Which the
Ketone Body D-β-Hydroxybutyrate
May Improve the Multiple Cellular
Pathologies of Parkinson's Disease
Front. Nutr. 6:63.
doi: 10.3389/fnut.2019.00063

Parkinson's disease, a progressive neurodegenerative disorder characterized by motor and non-motor symptoms, is strongly associated with the death of dopaminergic neurons in the brain's substantia nigra. Although dopamine replacement therapy temporarily helps patients manage their motor symptoms, this current standard of care fails to address the underlying network of pathologies that contribute to the persistent death of dopaminergic neurons. Thus, new treatment approaches are needed that address the underlying pathologies and, thereby, slow or halt the progression of the actual disease. D-β-hydroxybutyrate – a ketone body produced by the liver to support brain function during periods of starvation – may provide an option. Lifestyle interventions that induce endogenous D-β-hydroxybutyrate production, such as caloric restriction and ketogenic diets, are known to increase healthspan and lifespan in animal models and are used to treat neurological disorders. The efficacy of these ketosis-inducing interventions, along with the recent development of commercially available D-β-hydroxybutyrate-based nutritional supplements, should inspire interest in the possibility that D-β-hydroxybutyrate itself exerts neuroprotective effects. This review provides a molecular model to justify the further exploration of such a possibility. Herein, we explore the cellular mechanisms by which the ketone body, D-β-hydroxybutyrate, acting both as a metabolite and as a signaling molecule, could help to prevent the development, or slow the progression of, Parkinson's disease. Specifically, the metabolism of D-β-hydroxybutyrate may help neurons replenish their depleted ATP stores and protect neurons against oxidative damage. As a G-protein-coupled receptor ligand and histone deacetylase inhibitor, D-β-hydroxybutyrate may further protect neurons against energy deficit and oxidative stress, while also decreasing damaging neuroinflammation and death by apoptosis. Restricted to the available evidence, our model relies largely upon

the interpretation of data from the separate literatures on the cellular effects of D- β -hydroxybutyrate and on the pathogenesis of Parkinson's disease. Future studies are needed to reveal whether D- β -hydroxybutyrate actually has the potential to serve as an adjunctive nutritional therapy for Parkinson's disease.

Keywords: apoptosis, dopamine, D- β -hydroxybutyrate, energy metabolism, inflammation, oxidative stress, Parkinson's disease

INTRODUCTION

Parkinson's disease (PD) is the world's second most common and fastest growing neurodegenerative disorder. At present, PD affects 2–3% of individuals over the age of 65, a figure that is expected to double by the year 2040 (1, 2). Hence, if it were an infectious disease, PD would quite rightly be called an epidemic. Symptomatically, PD manifests in several classical motor symptoms, including tremors and bradykinesia, as well as in a wide variety of non-motor symptoms, such as disordered sleep and cognitive dysfunction. As there is no cure for PD, symptoms inevitably progress and inflict devastating consequences on individuals and on their families.

In common with Alzheimer and other neurodegenerative diseases, PD is biologically characterized by protein misfolding and the rampant death of neurons. Specifically, PD is characterized by the aggregation of α -synuclein protein and the death of dopaminergic neurons in the midbrain substantia nigra (SN), although PD affects other neurotransmitter systems as well. The pathological mechanisms underlying these biological hallmarks are highly complex and include both metabolic and signaling dysfunctions.


As both a fuel substrate and signaling molecule, the ketone body D- β -hydroxybutyrate (β HB) may be well-suited to slow, halt, or even reverse the progression of PD. Indeed, Tieu et al. have shown that β HB can successfully protect against the death of dopaminergic neurons in the SN and that β HB can alleviate the symptoms of PD in mice (3). At present, many publications are available describing either the ways in which β HB alters metabolism/cell signaling or how β HB affects specific pathological mechanisms that are associated with PD. This review attempts to synthesize these separate bodies of evidence in order to suggest a model for how β HB,


acting as both a metabolite and as a signaling molecule, might address many aspects of the pathological network underlying PD; and, therefore, that exogenous β HB supplementation may represent an improvement upon the current standard of care. In this review, we subcategorize β HB's effects on metabolism and cell signaling, respectively, into β HB's effects on (i) ATP production and (ii) antioxidant defenses, and into β HB's actions as (iii) an activator of the G-protein-coupled, hydroxycarboxylic acid receptor 2 (HCAR2) and as (iv) an inhibitor of histone deacetylases (HDACs). Throughout the review we attempt to identify how β HB, acting through these four strategies, can protect against the four most prominent pathological mechanisms in PD, namely abnormal cellular energy metabolism, oxidative stress, inflammation, and apoptosis (Figure 1).

ABNORMAL ENERGY METABOLISM

Due to their large size, extensive arborization and calcium-pacemaking activity, SN dopaminergic neurons are particularly metabolically active cells and are, therefore, especially susceptible

Abbreviations: ATGs, autophagy-related proteins; ATP, adenosine triphosphate; BDNF, brain-derived neurotrophic factor; BH₂, dihydrobiopterin; BH₄, tetrahydrobiopterin; β -HB, D- β -hydroxybutyrate; ER, endoplasmic reticulum; FOXO3A, forkhead box O3A; G6P, glucose-6-phosphate; HCAR2, hydroxycarboxylic acid receptor 2; HDAC, histone deacetylase; IL-1 β , interleukin-1 β ; iNOS, inducible nitric oxide synthase; mnSOD, manganese superoxide dismutase; MPTP, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; Mt2, metallothionein 2; NAD, nicotinamide adenine dinucleotide; NADP, nicotinamide adenine dinucleotide phosphate; NF κ B, nuclear factor κ -light-chain-enhancer of activated B cells; NO, nitric oxide; O₂ \cdot , superoxide; PARP1, poly ADP-ribose polymerase 1; PGC1- α , peroxisome proliferator-activated receptor gamma coactivator 1- α ; PD, Parkinson's disease; ³¹P-MRS, ³¹phosphorus-magnetic resonance spectroscopy; p53, tumor suppressor p53; Q, coenzyme Q; ROS, reactive oxygen species; Ru5P, ribulose-5-phosphate; SIRT1, silent mating type information regulation 2 homolog 1; SN, substantia nigra; TNF- α , tumor necrosis factor- α ; TRX, thioredoxin; ULK1, unc-51 like autophagy activating kinase 1.





to energy deprivation (1, 4). Indeed, ^{31}P -MRS imaging reveals that ATP levels in the region of the brain that includes the SN are significantly reduced in patients with PD (5). Evidence from mouse models suggests that this relationship is not simply correlative. MPTP, the complex I inhibitor most commonly used to produce animal models of PD, depletes cerebral ATP *in vivo* (6). Furthermore, pharmacologically blocking ATP consumption or increasing ATP production in PD mice is sufficient to prevent α -synuclein aggregation and the death of dopaminergic neurons in the SN, and is sufficient to protect against the motor symptoms of PD (7).

There are at least two mechanisms by which β HB may increase ATP levels in dopaminergic neurons. First, the work that launched contemporary scientific interest into the field of exogenous ketones, conducted by Sato et al. on perfused rat hearts, suggests that β HB metabolism increases mitochondrial ATP production by exerting opposite redox effects on the respiratory chain electron carriers, NAD and coenzyme Q (Q). By reducing (decreasing) the NAD^+/NADH ratio, while simultaneously oxidizing (increasing) the Q/QH_2 ratio, β HB increases the difference between the redox potentials of these two electron carrier couples (8). This increase in redox span is biochemically analogous to increasing the height span from which a bowling ball is dropped to the ground. In both cases, more energy is available to do work. Therefore, when electrons are passed down from NADH to Q , more protons can be pumped into the intermembrane space to drive the generation of more ATP by chemiosmosis. In this way, β HB metabolism can increase the redox span within the electron transport chain to increase the generation of ATP by oxidative phosphorylation (Figure 2).

Rodent data suggest that β HB metabolism also permits dopaminergic neurons to circumvent the blockade of complex I, a phenomenon that contributes to mitochondrial dysfunction


in human PD (9–11), by feeding electrons into the respiratory chain at complex II (Figure 2). This mechanism makes biochemical sense because the rate limiting step of β HB catabolism generates succinate, the oxidative fuel for complex II. Specifically, it has been shown that administration of β HB to MPTP-treated PD mice protects dopaminergic SN neurons from cell death and that this effect is blocked by the specific inhibition of complex II (3). In addition, β HB is able to increase ATP levels in brain mitochondria in the presence of MPTP-mediated complex I inhibition, but not when flux through complex I and complex II are both inhibited (3).

And, although flux through complex II can be linked to a decreased Q/QH_2 ratio, the redox span and complex II flux models are not contradictory because an increase in flux through a pathway does equate to an increase in the metabolites in that pathway. In fact, β HB only increases succinate levels when flux through complex II is blocked (3), a finding consistent with the notion that β HB can increase Q/QH_2 turnover without decreasing the ratio itself. Therefore, these two mechanisms, whereby β HB increases ATP production in PD, may be less contradictory than complimentary.

OXIDATIVE STRESS AND ANTIOXIDANTS

By altering the ratios of redox couples within mitochondria and the cytoplasm, β HB metabolism may both diminish the production of reactive oxygen species (ROS) and bolster antioxidant defenses. Most mitochondrial ROS are generated via the process of reverse electron transport, in which electrons are passed from QH_2 to oxygen at complex I to generate superoxide. Therefore, by oxidizing the Q/QH_2 couple (8), β HB not only increases the production of ATP (see above), but also decreases the production of ROS (Figure 3).

In addition to increasing the Q/QH_2 ratio and decreasing the NAD^+/NADH ratio (8), β HB also decreases the $\text{NADP}^+/\text{NADPH}$ ratio through a variety of mechanisms. First, NADH and NADPH are tightly linked, especially in neurons, by the direct transfer of hydride ions from NADH to NADP^+ via nicotinamide nucleotide transhydrogenase (12, 13). Second, by decreasing glycolytic flux, β HB forces glucose-6-phosphate down the pentose phosphate pathway, leading to the production of two equivalents of NADPH (8, 12–14). Third, by increasing ~ 15 -fold the concentration of mitochondrial acetyl-CoA (8), β HB increases the concentration of mitochondrial citrate and the export of this citrate into the cytoplasm by the citrate-pyruvate and citrate-isocitrate cycles, each of which includes an NADP^+ to NADPH reduction step catalyzed by malic enzyme and isocitrate dehydrogenase, respectively (12, 15). The products of these two carrier systems, oxaloacetate and α -ketoglutarate, are returned to the Krebs cycle to complete the circuit. Therefore, by increasing mitochondrial citrate concentrations, β HB forces the citrate-pyruvate and citrate-isocitrate wheels to spin faster to produce more NADPH (Figure 3). As all known intracellular antioxidants, either directly or indirectly, depend upon NADPH as the



electron donor, β HB-mediated reduction of NADP⁺ into NADPH translates into an increase in the reduced levels of glutathione, thioredoxin, vitamins C and E, and other essential antioxidants (15).

By increasing NADPH levels, β HB may exert another beneficial effect in the context of PD: β HB may increase the synthesis of dopamine and other neurotransmitters. As the ultimate reducing agent, NADPH supports the reduction of dihydrobiopterin into tetrahydrobiopterin, a coenzyme critical in the synthesis of dopamine, noradrenalin, serotonin, and melatonin. Although the possible effect of β HB on these neurotransmitter systems is speculative and requires further investigation, it is worth mentioning if for no other reason than this mechanism by which β HB might increase neurotransmitter levels is distinct from the current standard of care for PD, levodopa therapy, which only provides the direct precursor to dopamine. Therefore, β HB may be able to increase the synthesis, not only of dopamine (3), but also of other neurotransmitters that are underproduced in PD and which contribute to the non-motor symptoms of PD not addressed by levodopa therapy (15).

β HB not only increases the Q/QH₂ ratio to decrease the generation of ROS, and decreases the NADP⁺/NADPH ratio to increase antioxidant defenses and support neurotransmitter synthesis, but also has the unique ability to regulate all four of the “great” controlling nucleotide coenzyme couples: NAD⁺/NADH, NADP⁺/NADPH, acetyl-CoA/CoA, and ATP/ADP. These coenzymes diffuse throughout the cell, collectively regulating


at least 1,500 known enzymatic reactions through their shared potential energies. Therefore, by regulating these four couples, β HB metabolism shapes the entire metabolome in ways that we are only beginning to appreciate and may affect not only energy crisis and oxidative stress, but also inflammation and apoptosis (15).

G-PROTEIN-COUPLED RECEPTOR (HCAR2) ACTIVATION

In addition to its role as a fuel substrate, β HB is a ligand for hydroxycarboxylic acid receptor 2 (HCAR2), a G-protein coupled receptor that is upregulated in the SN of PD patients (14, 16, 17). Although HCAR2 is upregulated in the PD brain, it is predicted to be underactive because PD patients also exhibit lower levels niacin, another HCAR2 ligand (17). Therefore, by substituting for niacin and activating upregulated HCAR2, β HB may be able to target a set of sensitized pathways in the PD brain, pathways that include the critical proteins SIRT1 and NF κ B (Figure 4).

SIRT1 is a deacetylase whose activity is strongly associated with the generic health and longevity benefits of caloric restriction and that is underactive in the PD brain (18, 19). One way in which SIRT1 is thought to mediate its neurological benefits is by upregulating autophagy, a catabolic process active in all cells that facilitates the degradation and recycling of damaged cellular components. SIRT1 activates a handful of autophagy proteins, including ULK1 and several ATGs, to increase the activity of this essential cellular recycling process and promote the clearance of defective and damaged mitochondria (20, 21). In a complementary fashion, SIRT1 activates the master regulator of mitochondrial biogenesis, PGC1- α , which is also underexpressed in, and has been strongly implicated in the pathogenesis of, PD (20, 22–24). Not only can SIRT1 increase autophagy and mitochondrial biogenesis, but it also induces FOXO3A-dependent expression of the antioxidant genes catalase, mnSOD, and Mt2 (20). Therefore, the stimulation of SIRT1 by β HB could protect cells against energy depletion and oxidative stress by improving the health of the mitochondrial pool and by bolstering antioxidant defenses. In addition, rodent studies suggest that a ketogenic diet induces SIRT1-dependent deacetylation of the transcription factor p53 and, thereby, increases the expression of anti-apoptotic proteins, decreases the expression of pro-apoptotic proteins, and ultimately protects neurons from apoptosis (25) (Figure 4).

By binding to HCAR2 on macrophages and microglia in the brain, β HB also inhibits NF κ B-mediated neuroinflammation, a critical pathological feature in PD (26–28) (Figure 4). NF κ B is a potent proinflammatory transcription factor that is elevated in the PD brain (29), has been proposed as a therapeutic target for PD (30), and induces the expression of TNF- α , IL-1 β , and inducible nitric oxide synthase (iNOS) (31–33). In addition to stimulating an innate immune response in the brain, the cytokines TNF- α and IL-1 β , which have likewise been proposed as targets for potential PD treatments (34),




can also directly promote apoptosis by binding to death receptors on neurons (35, 36). The NO generated by iNOS, which is also pathologically important in PD (37), is likewise a proinflammatory molecule that exerts other pathological influences. NO can block glutamate reuptake, leading to excitotoxicity and apoptosis (32). NO can also provoke oxidative stress by post-translationally modifying proteins important in mitochondrial quality control and can induce nitroxidative stress by combining with superoxide to form peroxynitrite and other cytotoxic molecules (32).

HISTONE DEACETYLASE INHIBITION

It has been shown *in vitro* that histone deacetylase (HDAC) inhibition decreases α -synuclein toxicity and protects dopaminergic neurons from cell death (38). As a natural inhibitor of HDACs 1, 3, and 4 (16, 39), β HB may manage each of the pathologies underlying PD by regulating HDACs and altering gene expression.

For example, β HB-mediated HDAC inhibition in mice increases the expression of brain-derived neurotrophic factor (BDNF) (40), a molecule renowned for its putative ability to stimulate adult neurogenesis. In patients with PD, BDNF expression in the SN is significantly decreased (41), and, in rodent and primate models of PD, BDNF has been shown to protect dopaminergic SN neurons from cell death (42, 43). In mouse primary cortical neurons, β HB-mediated HDAC inhibition increases BDNF expression, decreases the NAD⁺/NADH ratio, and increases ATP (40, 44). BDNF also prevents NF κ B-mediated neuroinflammation and apoptosis



in rodent models of central nervous system inflammatory diseases (45, 46). Finally, BDNF increases the activity of multiple antioxidant enzymes and decreases oxidative damage in the basal ganglia of PD rats (47, 48). Therefore, by increasing BDNF expression alone, β HB may protect against energy depletion, oxidative stress, inflammation, and apoptosis (Figure 5).

BDNF is one representative example of how β HB-mediated HDAC inhibition can exert a wide range of neuroprotective effects by altering transcription; another example being β HB-mediated upregulation of the antioxidant genes catalase, mnSOD, and Mt2 (16, 49) (Figure 5). In fact, the β HB target, HDAC4, is responsible for the characteristic pattern of gene expression that differentiates PD neurons from healthy neurons (39). Furthermore, HDAC4 inhibition rescues gene expression in dopaminergic neurons derived from patients with PD. This genetic rescue, in turn, decreases endoplasmic reticulum stress, which is a cellular response to protein aggregation marked by (i) oxidative stress, (ii) apoptosis (50, 51), and (iii) glucocerebrosidase degradation (a critical autophagy-related enzyme already decreased in the PD brain) (52, 53) (Figure 5).


CONCLUDING REMARKS

Segregating the pathologies underlying PD is, to some extent, an arbitrary exercise because they are deeply interconnected. Energetic abnormalities cause cells, which have been unable to maintain homeostasis, and thus may have compromised genomes, to undergo apoptosis (54). Apoptosis can deplete the extracellular pool of neuroprotective BDNF, thereby increasing inflammation (45, 46). The NO produced as part of the

inflammatory response can induce oxidative stress (32), which can cause mitochondrial damage and increase the energy crisis (Figure 6, outermost loop).

Of course, PD pathology is not a simple loop, but a complex network. Oxidative stress induces DNA damage, exciting the activity of the DNA repair protein, PARP1, which catabolizes the SIRT1 cofactor, NAD^+ (25), and thus precipitates further oxidative stress, energy crisis, and apoptosis. ROS also directly induces apoptosis and NF κ B-mediated inflammation, the latter of which accentuates oxidative stress, causes apoptosis, and, because NF κ B is induced by the cytokines that it induces, more inflammation (31) (Figure 6).

The pathologies that underlay PD are synergistic. This interconnectedness undermines any intervention aimed at a single point within the network. Perhaps that is why no true preventative treatments exist and why the current standard of care remains the largely palliative option of dopamine replacement therapy. However, there is a flipside to this complexity, for any molecule that can target many points within the network, as β HB can, may capitalize on the feedback loops, such that the multiple mechanisms by which it operates add up to more than the sum of their parts. In this review, we attempted to integrate the still largely independent bodies of literature on the biochemical effects of β HB and on that pathological mechanisms of PD in order to present a model



for how β HB could be used to improve the multiple cellular pathologies of PD. This model, in combination with preliminary data that show the ketogenic diet may be ameliorative in PD (55), suggests that exogenous β HB, now available as an FDA-approved sports drink, may have therapeutic potential for the prevention and/or treatment of PD. Future studies, including those currently being conducted, will reveal the authenticity of this potential.

REFERENCES

- Poewe W, Seppi K, Tanner CM, Halliday GM, Brundin P, Volkman J, et al. Parkinson disease. *Nat Rev Dis Primers*. (2017) 3:17013. doi: 10.1038/nrdp.2017.13
- Dorsey ER, Bloem BR. The Parkinson pandemic - a call to action. *JAMA Neurol*. (2018) 75:9–10. doi: 10.1001/jamaneurol.2017.3299
- Tieu K, Perier C, Caspersen C, Teismann P, Wu DC, Yan SD, et al. D- β -Hydroxybutyrate rescues mitochondrial respiration and mitigates features of Parkinson disease. *J Clin Invest*. (2003) 112:892–901. doi: 10.1093/brinp/123.2.340
- Dias V, Junn E, Mouradian MM. The role of oxidative stress in Parkinson's disease. *J Parkinsons Dis*. (2013) 3:461–91. doi: 10.3233/JPD-130230
- Hu MT, Taylor-Robinson SD, Chaudhuri KR, Bell JD, Labbé C, Cunningham VJ, et al. Cortical dysfunction in non-demented Parkinson's disease patients - a combined 31 P-MRS and 18 FDG-PET study. *Brain*. (2000) 123:340–52. doi: 10.1093/brain/123.2.340
- Chan P, DeLanney LE, Irwin I, Langston JW, Di Monte D. Rapid ATP loss caused by 1-methyl-4-phenyl-1,2,3,6 in mouse brain. *J Neurochem*. (1991) 57:348–83. doi: 10.1111/j.1471-4159.1991.tb02134.x
- Nakano M, Imamura H, Sasaoka N, Yamamoto M, Uemura N, Shudo T, et al. ATP maintenance via two types of ATP regulators mitigates pathological phenotypes in mouse models of Parkinson's disease. *EBioMedicine*. (2017) 22:225–41. doi: 10.1016/j.ebiom.2017.07.024
- Sato K, Kashiwaya Y, Keon CA, Tsuchiya N, King MT, Radda GK, et al. Insulin, ketone bodies, and mitochondrial energy transduction. *FASEB J*. (1995) 9:651–8. doi: 10.1096/fasebj.9.8.7768357
- Mann VM, Cooper JM, Krige D, Daniel SE, Schapira AH, Marsden CD. Brain, skeletal muscle and platelet homogenate mitochondrial function in Parkinson's disease. *Brain*. (1992) 115:333–42. doi: 10.1093/brain/115.2.333
- Devi L, Raghavendran V, Prabhu BM, Avadhani NG, Anandatheerthavarada HK. Mitochondrial import and accumulation of α -synuclein impair complex I in human dopaminergic neuronal cultures and Parkinson disease brain. *J Biol Chem*. (2008) 283:9089–100. doi: 10.1074/jbc.M710012200
- Benecke R, Strümpfer P, Weiss H. Electron transfer complexes I and IV of platelets are abnormal in Parkinson's disease but normal in Parkinson-plus syndromes. *Brain*. (1993) 116:1451–63. doi: 10.1093/brain/116.6.1451
- Agedal L, Niere M, Ziegler M. The phosphate makes a difference: cellular functions of NADP. *Redox Rep*. (2010) 15:2–10. doi: 10.1179/174329210X12650506623122
- Blacker TS, Duchon MR. Investigating mitochondrial redox state using NADH and NADPH autofluorescence. *Free Radic Biol Med*. (2016) 100:53–65. doi: 10.1016/j.freeradbiomed.2016.08.010
- Veech RL, Bradshaw PC, Clarke K, Curtis W, Pawlosky R, Todd King M. Ketone bodies mimic the life span extending properties of caloric restriction. *IUBMB Life*. (2017) 69:305–14. doi: 10.1002/iub.1627
- Veech RL, Todd King M, Pawlosky R, Kashiwaya Y, Bradshaw PC, Curtis W. The 'great' controlling nucleotide coenzymes. *IUBMB Life*. (2019) 71:565–79. doi: 10.1002/iub.1997
- Newman J, Verdin E. Ketone bodies as signaling metabolites. *Trends Endocrinol Metab*. (2014) 25:42–52. doi: 10.1016/j.tem.2013.09.002
- Wakade C, Chong R, Bradley E, Thomas B, Morgan J. Upregulation of GPR109A in Parkinson's disease. *PLoS ONE*. (2014) 9:e109818. doi: 10.1371/journal.pone.0109818

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

FUNDING

NN would like to gratefully acknowledge the Keasbey Memorial Foundation for funding his research at Oxford.

- Rahman S, Islam R. Mammalian Sirt1: insights on its biological functions. *Cell Commun Signal*. (2011) 9:11. doi: 10.1186/1478-811X-9-11
- Singh P, Hanson PS, Morris CM. SIRT1 ameliorates oxidative stress induced neural cell death and is down-regulated in Parkinson's disease. *BMC Neurosci*. (2017) 18:46. doi: 10.1186/s12868-017-0364-1
- McCarty ME, DiNicolantonio JJ, O'Keefe JH. Ketosis may promote brain macroautophagy by activating Sirt1 and hypoxia-inducible factor-1. *Med Hypotheses*. (2015) 85:631–9. doi: 10.1016/j.mehy.2015.08.002
- Herzig S, Shaw RJ. AMPK: guardian of metabolism and mitochondrial homeostasis. *Nat Rev Mol Cell Biol*. (2018) 19:121–35. doi: 10.1038/nrm.2017.95
- Ciron C, Zheng L, Bobela W, Knott GW, Leone TC, Kelly DP, et al. PGC-1 α activity in nigral dopamine neurons determines vulnerability to α -synuclein. *Acta Neuropathol Commun*. (2015) 3:16. doi: 10.1186/s40478-015-0200-8
- Su X, Chu Y, Kordower JH, Li B, Cao H, Huang L, et al. PGC-1 α promoter methylation in Parkinson's disease. *PLoS ONE*. (2015) 10:24. doi: 10.1371/journal.pone.0134087
- Zheng B, Liao Z, Locascio JJ, Lesniak KA, Roderick SS, Watt ML, et al. PGC-1, a potential therapeutic target for early intervention in Parkinson's disease. *Sci Transl Med*. (2010) 2:52ra73. doi: 10.1126/scitranslmed.3001059
- Scheibye-Knudsen M, Mitchell SJ, Fang EF, Iyama T, Ward T, Wang J, et al. A high-fat diet and NAD⁺ activate Sirt1 to rescue premature aging in cockayne syndrome. *Cell Metab*. (2014) 20:840–55. doi: 10.1016/j.cmet.2014.10.005
- Rahman H, Muhammad S, Khan MA, Chen H, Ridder DA, Müller-Fielitz H, et al. The β -hydroxybutyrate receptor HCA 2 activates a neuroprotective subset of macrophages. *Nat Commun*. (2014) 5:3944. doi: 10.1038/ncomms4944
- Parodi B, Rossi S, Morando S, Cordano C, Bragioni A, Motta C, et al. Fumarates modulate microglia activation through a novel HCAR2 signaling pathway and rescue synaptic dysregulation in inflamed CNS. *Acta Neuropathol*. (2015) 130:279–95. doi: 10.1007/s00401-015-1422-3
- Troncoso-Escudero P, Parra A, Nassif M, Vidal RL. Outside in: unraveling the role of neuroinflammation in the progression of Parkinson's disease. *Front Neurol*. (2018) 9:860. doi: 10.3389/fneur.2018.00860
- Hunot S, Brugg B, Ricard D, Michel PP, Muriel M-P, Ruberg M, et al. Nuclear translocation of NF- κ B is increased in dopaminergic neurons of patients with Parkinson disease. *Proc Natl Acad Sci USA*. (1997) 94:7531–6. doi: 10.1073/pnas.94.14.7531
- Flood PM, Qian L, Peterson LJ, Zhang F, Shi JS, Gao HM, et al. Transcriptional factor NF- κ B as a target for therapy in Parkinson's disease. *Parkinsons Dis*. (2011) 2011:216298. doi: 10.4061/2011/216298
- Shabab T, Khanabdali R, Moghadamtousi SZ, Kadir HA, Mohan G. Neuroinflammation pathways: a general review. *Int J Neurosci*. (2017) 127:624–33. doi: 10.1080/00207454.2016.1212854
- Yuste JE, Tarragon E, Campuzano CM, Ros-Bernal F. Implications of glial nitric oxide in neurodegenerative diseases. *Front Cell Neurosci*. (2015) 9:e1–13. doi: 10.3389/fncel.2015.00322
- Digby JE, Martinez F, Jefferson A, Ruparelia N, Chai J, Wamil M, et al. Anti-inflammatory effects of nicotinic acid in human monocytes are mediated by GPR109A dependent mechanisms. *Arterioscler Thromb Vasc Biol*. (2012) 32:669–76. doi: 10.1161/ATVBAHA
- Leal MC, Casabona JC, Puntel M, Pitossi FJ. Interleukin-1 β and tumor necrosis factor- α : reliable targets for protective therapies in Parkinson's disease? *Front Cell Neurosci*. (2013) 7:53. doi: 10.3389/fncel.2013.00053

35. Taylor RC, Cullen SP, Martin SJ. Apoptosis: controlled demolition at the cellular level. *Nat Rev Mol Cell Biol.* (2008) 9:231–41. doi: 10.1038/nrm2312
36. Cui J, Zhang M, Zhang YQ, Xu ZH. JNK pathway: diseases and therapeutic potential. *Acta Pharmacol Sin.* (2007) 28:601–8. doi: 10.1111/j.1745-7254.2007.00579.x
37. Zhang L, Dawson VL, Dawson TM. Role of nitric oxide in Parkinson's disease. *Pharmacol Ther.* (2006) 109:33–41. doi: 10.1016/j.pharmthera.2005.05.007
38. Kontopoulos E, Parvin JD, Feany MB. α -synuclein acts in the nucleus to inhibit histone acetylation and promote neurotoxicity. *Hum Mol Genet.* (2006) 15:3012–23. doi: 10.1093/hmg/ddl243
39. Lang C, Campbell KR, Ryan BJ, Carling P, Attar M, Vowles J, et al. Single-cell sequencing of iPSC-dopamine neurons reconstructs disease progression and identifies HDAC4 as a regulator of Parkinson cell phenotypes. *Cell Stem Cell.* (2019) 24:93–106. doi: 10.1016/j.stem.2018.10.023
40. Sleiman SF, Henry J, Al-Haddad R, El Hayek L, Abou Haidar E, Stringer T, et al. Exercise promotes the expression of Brain Derived Neurotrophic Factor (BDNF) through the action of the ketone body β -hydroxybutyrate. *Elife.* (2016) 5:e15092. doi: 10.7554/eLife.15092
41. Howells DW, Porritt MJ, Wong JY, Batchelor PE, Kalnins R, Hughes AJ, et al. Reduced BDNF mRNA expression in the Parkinson's disease substantia nigra. *Exp Neurol.* (2000) 166:127–35. doi: 10.1006/exnr.2000.7483
42. Frim DM, Uhler TA, Galpern WR, Beal MF, Breakefield XO, Isacson O. Implanted fibroblasts genetically engineered to produce brain-derived neurotrophic factor prevent 1-methyl-4-phenylpyridinium toxicity to dopaminergic neurons in the rat. *Proc Natl Acad Sci USA.* (1994) 91:5104–8. doi: 10.1073/pnas.91.11.5104
43. Tsukahara T, Takeda M, Shimohama S, Ohara O, Hashimoto N. Effects of drain-derived neurotrophic factor on 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced Parkinsonism in monkeys. *Neurosurgery.* (1995) 37:733–41. doi: 10.1097/00006123-199510000-00018
44. Marosi K, Kim SW, Moehl K, Scheibye-Knudsen M, Cheng A, Cutler R, et al. 3-Hydroxybutyrate regulates energy metabolism and induces BDNF expression in cerebral cortical neurons. *J Neurochem.* (2016) 139:769–81. doi: 10.1111/jnc.13868
45. Xu D, Lian D, Wu J, Liu Y, Zhu M, Sun J, et al. Brain-derived neurotrophic factor reduces inflammation and hippocampal apoptosis in experimental *Streptococcus pneumoniae* meningitis. *J Neuroinflammation.* (2017) 14:156. doi: 10.1186/s12974-017-0930-6
46. Makar TK, Trisler D, Sura KT, Sultana S, Patel N, Bever CT. Brain derived neurotrophic factor treatment reduces inflammation and apoptosis in experimental allergic encephalomyelitis. *J Neurol Sci.* (2008) 270:70–6. doi: 10.1016/j.jns.2008.02.011
47. Mattson MP, Lovell MA, Furukawa K, Markesbery WR. Neurotrophic factors attenuate glutamate-induced accumulation of peroxides, elevation of intracellular Ca^{2+} concentration, and neurotoxicity and increase antioxidant enzyme activities in hippocampal neurons. *J Neurochem.* (1995) 65:1740–51. doi: 10.1046/j.1471-4159.1995.65041740.x
48. Kirschner PB, Jenkins BG, Schulz JB, Finkelstein SP, Matthews RT, Rosen BR, et al. NGF, BDNF and NT-5, but not NT-3 protect against MPP+ toxicity and oxidative stress in neonatal animals. *Brain Res.* (1996) 713:178–85. doi: 10.1016/0006-8993(95)01513-2
49. Shimazu T, Hirschev MD, Newman J, He W, Shirakawa K, Le Moan N, et al. Suppression of oxidative stress by β -hydroxybutyrate, an endogenous histone deacetylase inhibitor. *Science.* (2013) 339:211–4. doi: 10.1126/science.1227166
50. Zhao L, Ackerman S. Endoplasmic reticulum stress in health and disease. *Curr Opin Cell Biol.* (2006) 18:444–52. doi: 10.1016/j.ceb.2006.06.005
51. Lai E, Teodoro T, Volchuk A. Endoplasmic reticulum stress: signaling the unfolded protein response. *Physiology.* (2007) 22:193–201. doi: 10.1152/physiol.00050.2006
52. Gegg ME, Burke D, Heales SJ, Cooper JM, Hardy J, Wood NW. Glucocerebrosidase deficiency in substantia nigra of parkinson disease brains. *Ann Neurol.* (2012) 72:455–63. doi: 10.1002/ana.23614
53. Schapira AHV, Gegg ME. Glucocerebrosidase in the pathogenesis and treatment of Parkinson disease. *Proc Natl Acad Sci USA.* (2013) 110:3214–5. doi: 10.1073/pnas.1300822110
54. Izyumov DS, Avetisyan AV, Pletjushkina OY, Sakharov DV, Wirtz KW, Chernyak BV, et al. 'Wages of Fear': transient threefold decrease in intracellular ATP level imposes apoptosis. *Biochim Biophys Acta Bioenerg.* (2004) 1658:141–7. doi: 10.1016/j.bbabi.2004.05.007
55. Vanitallie TB, Nonas C, Di Rocco A, Boyar K, Hyams K, Heymsfield SB. Treatment of Parkinson disease with diet-induced hyperketonemia: a feasibility study. *Neurology.* (2005) 64:728–30. doi: 10.1212/01.WNL.0000152046.1390.45

Conflict of Interest Statement: Intellectual property covering uses of dietary ketone and ketone ester supplementation is owned by BTG Ltd., the University of Oxford, the National Institute of Health and TΔS Ltd. Should royalties ever accrue from these patents, KC, as inventor, will receive a share of the royalties under the terms prescribed by the University of Oxford. KC is a director of TΔS Ltd., a company spun out of the University of Oxford to develop and commercialize products based on the science of ketone bodies in human nutrition.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Norwitz, Hu and Clarke. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.